

Automorphism covariant representations of the holonomy-flux $\ast$ -algebra

Andrzej Okołów^{1*} and Jerzy Lewandowski^{1,2†}

June 14, 2003

1. *Instytut Fizyki Teoretycznej, Uniwersytet Warszawski, ul. Hoża 69,
00-681 Warsaw, Poland*
2. *Center for Gravitational Physics and Geometry, Physics Department,
104 Davey, Penn State, University Park, PA 16802, USA*

* *oko@fuw.edu.pl*

† *lewand@fuw.edu.pl*

Abstract

We continue the analysis of representations of cylindrical functions and fluxes which are commonly used as elementary variables of Loop Quantum Gravity. We consider an arbitrary principal bundle of a compact connected structure group and, following Sahlmann's ideas [1], define a holonomy-flux $\ast$ -algebra whose elements correspond to the elementary variables. There exists a natural action of automorphisms of the bundle on the algebra; this action generalizes the action of analytic diffeomorphisms and gauge transformations on the algebra considered in earlier works. We define the automorphism covariance of a $\ast$ -representation of the algebra on a Hilbert space and prove that the only Hilbert space admitting such a representation is a direct sum of the spaces L^2 , given by a unique measure on the space of generalized connections. This result is a generalization of our previous work [2] where we assumed that the principal bundle is trivial and its base manifold is $\mathbb{R}^d$.

1 Introduction

Loop Quantum Gravity¹ (LQG) is an attempt to obtain a consistent theory of quantum gravity quantizing the classical theory of general relativity (GR).

¹For a comprehensive introduction and references to the subject see [3].

The departure point of LQG is a Hamiltonian formulation of GR whose configuration space is the space of connections on an $SU(2)$ principal fibre bundle over three-dimensional 'spatial' manifold. The passage from the classical theory to the quantum theory is done by canonical quantization which is required to preserve symmetries of the Hamiltonian formulation of GR. This means, in particular, that the quantization should not make use of any geometrical background structure on the manifold, e.g. a fixed metric.

A crucial step of the canonical quantization procedure is the choice of elementary variables which are (complex) functions on the phase space of the theory (see e.g. [4]). Once such a set is chosen, the quantization procedure consists in assigning to any elementary variable an operator on a Hilbert space in such a way that (i) $(i\hbar)^{-1}$ times the commutator of the two operators assigned to a pair of variables corresponds to the Poisson bracket of these variables and (ii) the conjugate of the operator assigned to a variable corresponds to the complex conjugate of this variable — if $\hat{f}, \hat{g}$ are the operators assigned respectively to variables f, g then

$$(i) \quad (i\hbar)^{-1}[\hat{f}, \hat{g}] = \widehat{\{f, g\}}; \quad (ii) \quad \hat{f}^* = \widehat{\bar{f}}.$$

The assignment $f \mapsto \hat{f}$ is called a representation of elementary variables.

The present paper concerns the theory of representations of the commonly used elementary variables of LQG which are cylindrical functions and fluxes. Each (classical) cylindrical function depends on the holonomies of the $SU(2)$ -connections along a finite number of paths and each (classical) flux is given by an integral of the momentum variable canonically conjugate to the connection over a surface embedded in the 'spatial' manifold (the momentum can be naturally viewed as a differential two-form) [5, 6] (for a comprehensive description of the algebra of elementary variables see [7]). Because of the gauge and diffeomorphism invariance of LQG any admissible representation of the variables has to be *covariant* (in an appropriate sense) with respect to the gauge transformations and the action of diffeomorphisms. By now one knows only one non-trivial admissible representation of the variables, hence there arises a question about the *uniqueness* of the representation.

The question was addressed for the first time by Sahlmann [1, 8] who also proposed a new, more algebraic point of view on the issue. In Sahlmann's approach one defines a $*$ -algebra $\mathfrak{A}$ called a holonomy-flux $*$ -algebra whose elements correspond to cylindrical functions and fluxes, and then studies $*$ -representations of the algebra on a Hilbert space. The space of cylindrical functions used to construct $\mathfrak{A}$ can be completed to a C^* -algebra Cyl whose

Gelfand spectrum is naturally viewed as the space $\overline{\mathcal{A}}$ of so called generalized connections [5, 9]. The first result by Sahlmann is that each representation of the holonomy-flux $*$ -algebra on a Hilbert space induces a family of measures on $\overline{\mathcal{A}}$ [1]. In [8] Sahlmann studied the diffeomorphism covariance of $*$ -representations of the algebra $\mathfrak{A}$ built over $U(1)$ -connections. He proved that *every measure given by such a representation coincides with the measure introduced on $\overline{\mathcal{A}}$ by Ashtekar and Lewandowski* [10]. This result was generalized in [11] and [2] to the case of a holonomy-flux $*$ -algebra built over connections with any compact connected structure group.

In our earlier work [2], we assumed that the holonomy-flux $*$ -algebra is built over a *trivial* principal bundle over $\mathbb{R}^d$ ($d \geq 2$) as a base ('spatial') manifold. The goal of the present paper is to generalize the results of [2] to the case of a not necessarily trivializable bundle over an arbitrary base manifold. Thus in the sequel we will consider a principal bundle $P(\Sigma, G)$ such that (i) the base ('spatial') manifold Σ is real-analytic² and its dimension d is greater than 1 and (ii) the structure Lie group G is compact and connected. We will define a holonomy-flux $*$ -algebra $\mathfrak{A}$ associated with such a bundle and will call it the Sahlmann algebra. Next, we will have to define an action of diffeomorphisms of Σ and gauge transformations on the algebra. However, in general there is no canonical action of the diffeomorphisms on the space of connections and, consequently, on the algebra $\mathfrak{A}$. Thus instead of diffeomorphisms and gauge transformations treated separately we will use *automorphisms* of the bundle which unify the notion of diffeomorphisms and gauge transformations. Consequently, we will consider a natural action of the automorphisms on $\mathfrak{A}$ and define the notion of the *automorphism covariance* of a $*$ -representation of $\mathfrak{A}$ on a Hilbert space (precisely speaking, we will consider only those smooth automorphisms of $P(\Sigma, G)$ which define *analytic* diffeomorphisms on the base manifold Σ). As a main result of the paper, we will show that the only measure on $\overline{\mathcal{A}}$ which admits an automorphism covariant $*$ -representation of the Sahlmann algebra is the Ashtekar-Lewandowski measure.

In this paper we are able to prove merely the uniqueness of the mea-

²So far we are not able to construct a holonomy-flux $*$ -algebra in the case of any smooth base manifold Σ . The reason is that the operators generating the algebra are associated with some curves in Σ and with some $(d - 1)$ -dimensional submanifolds of Σ ($d = \dim \Sigma$) and the composition of the operators, i.e. the algebra multiplication, depends on the way in which a given curve intersects a given submanifold. Smooth curves and smooth submanifolds (unlike analytic ones) can intersect each other in a very complicated way which is an obstacle for defining the algebra multiplication in the case of any smooth manifold.

sure on $\overline{\mathcal{A}}$ which admits an automorphism covariant representation of the Sahlmann algebra, while the question of the uniqueness of the representation was left open. However, one can go a step further and prove the uniqueness of an automorphism covariant representation of the algebra which will be shown in [12].

Let us finally emphasize that in the present paper we will skip the 'derivation' of the Sahlmann algebra from the elementary variables, i.e. we will just give the definition of the algebra. The reader can find such a derivation, well suited for the considerations presented in the sequel, in [2].

2 Preliminaries

Let Σ be a real-analytic d -dimensional manifold ($d \geq 2$) and let $P(\Sigma, G)$ be a principal bundle over Σ , whose structure Lie group G is compact and connected. The right action of the group on P will be denoted by R_g :

$$P \times G \ni (p, g) \mapsto R_g(p) \in P,$$

and pr will denote the projection from P onto Σ defined by R_g .

Let M be any (possibly 0-dimensional) submanifold in Σ and let P_M denote the restriction of the bundle P to M . We say that a map $\theta : P_M \rightarrow P$ is a smooth morphism on P_M if (i) θ is an invertible smooth map and θ^{-1} is smooth and (ii) θ commutes with the right action R_g for every $g \in G$. The morphism θ is an automorphism of a bundle P_M if $\theta(P_M) = P_M$. Denote by Θ the set of all smooth automorphisms of P .

Every automorphism $\theta \in \Theta$ defines a smooth diffeomorphism $\check{\theta}$ on Σ such that:

$$\check{\theta} \circ \text{pr} = \text{pr} \circ \theta.$$

The automorphism θ is *vertical* if and only if $\check{\theta} = \text{id}$. Clearly, the set of all vertical automorphisms is a subgroup of Θ . Another subgroup of Θ is the one containing all automorphisms such that they define *analytic* diffeomorphisms on Σ :

$$\Theta^\omega := \{\theta \mid \theta \in \Theta \text{ and } \check{\theta} \text{ is analytic}\}.$$

Given an automorphism $\theta \in \Theta$ and a morphism θ' on P_M we define a morphism $\text{ad}_\theta \theta'$ on $P_{\check{\theta}(M)}$ in the following way:

$$P_{\check{\theta}(M)} \ni p \mapsto \text{ad}_\theta(\theta'(p)) := \theta(\theta'(\theta^{-1}(p))) \in P. \quad (2.1)$$

Let $\mathcal{A}$ be a set of all smooth connections on P . We will describe every element of $\mathcal{A}$ by means of a connection one-form A on P valued in the Lie

algebra G' of the group G . Usually, one describes a connection by a (family of) one-form(s) on the base manifold Σ , which is the pull-back of A with respect to a (family of local) section(s) of P . However, an arbitrary bundle P does not possess any distinguished section—therefore we will avoid the latter way of describing connections.

3 Sahlmann algebra

The Sahlmann algebra is an algebra of operators acting on a linear space of some complex functions defined on $\mathcal{A}$. Operators constituting the algebra are associated with some submanifolds embedded in Σ . We impose some regularity conditions on the submanifolds underlying the algebra and call them regular analytic submanifolds [2]:

Definition 3.1 (i) *0-dimensional regular analytic submanifold of Σ is a one-element subset of Σ ; (ii) an m -dimensional ($0 < m < \dim \Sigma$) regular analytic submanifold M is an analytic orientable m -dimensional submanifold such that its closure $\overline{M}$ is a compact subset of an analytic m -dimensional submanifold of Σ and $\overline{M} \setminus M$ is a finite union of regular analytic submanifolds of lower dimensions.*

The following subsets of Σ will serve to define the space of functions, on which Sahlmann algebra acts.

Definition 3.2 (i) *An analytic oriented edge in Σ is the closure of any connected 1-dimensional regular analytic submanifold with fixed orientation. (ii) An analytically embedded graph γ in Σ is a finite set of analytic oriented edges in Σ such that two distinct edges can intersect each other only at their endpoints. The edges will be called the edges of the graph γ , and their endpoints — the vertices of γ .*

3.1 Cylindrical functions

In this section we will introduce the notion of cylindrical function following [13].

Let e be an oriented edge. The orientation of e allows us to call one of its endpoints the source and the other one—the target of e . We will denote them by $(e)_s$ and $(e)_t$ respectively. Each connection A defines a parallel transport $A(e)$ along e being an isomorphism from $P_{(e)_s}$ onto $P_{(e)_t}$, which

commutes with the right action R_g for every $g \in G$; this means that $A(e)$ is a morphism on $P_{(e)_s}$ of image $P_{(e)_t}$. We also have:

$$A(e_1 \circ e_2) = A(e_1) \circ A(e_2), \quad A(e^{-1}) = A^{-1}(e), \quad (3.1)$$

where $e_1 \circ e_2$ denotes the composition of two edges such that $(e_1)_s = (e_2)_t$ and e^{-1} (the inverse of e) is an edge obtained from e by the change of the orientation.

Consider now a graph γ constituted by edges $\{e_1, \dots, e_N\}$ and define an equivalence relation on $\mathcal{A}$ in the following way. Given $A, A' \in \mathcal{A}$

$$A \sim_\gamma A' \text{ if and only if } A(e_I) = A'(e_I) \quad (3.2)$$

for every $I \in \{1, \dots, N\}$. Denote by A_γ the equivalence class of a connection A and by $\mathcal{A}_\gamma$ —the set of all equivalence classes defined by the relation. The relation $\sim_\gamma$ defines the following projection:

$$\mathcal{A} \ni A \mapsto p_\gamma(A) := A_\gamma \in \mathcal{A}_\gamma.$$

An important fact is that graphs in Σ form a directed set with relation $\geq$ defined as follows: $\gamma' \geq \gamma$ if and only if (i) every edge of γ can be expressed as a composition of edges of γ' and their inverses and (ii) each vertex of γ is a vertex of γ' [9]. It follows from (3.1) that if $\gamma' \geq \gamma$ then there exists a surjective projection $p_{\gamma\gamma'} : \mathcal{A}_{\gamma'} \rightarrow \mathcal{A}_\gamma$ such that:

$$p_\gamma = p_{\gamma\gamma'} \circ p_{\gamma'}. \quad (3.3)$$

Let $P_{V(\gamma)}$ be the bundle obtained by the restriction of the bundle P to the set $V(\gamma)$ of the vertices of γ . Then every A_γ can be thought as an N -tuple of morphisms

$$(A(e_1), \dots, A(e_N)) = (A(e_I))$$

acting on appropriate fibres of $P_{V(\gamma)}$. The bundle $P_{V(\gamma)}$ is trivializable; every trivialization of $P_{V(\gamma)}$ defines a bijection from $\mathcal{A}_\gamma$ onto G^N [10] and hence a structure of a (real-analytic) manifold on $\mathcal{A}_\gamma$. It turns out that this structure does not depend on the choice of the trivialization. The projection $p_{\gamma\gamma'}$ mentioned above is an analytic map from $\mathcal{A}_{\gamma'}$ onto $\mathcal{A}_\gamma$.

Definition 3.3 *We say that $\Psi : \mathcal{A} \rightarrow \mathbb{C}$ is a cylindrical function compatible with a graph γ , iff:*

$$\Psi = p_\gamma^* \psi. \quad (3.4)$$

where ψ is a complex function on $\mathcal{A}_\gamma$.

Each cylindrical function is compatible with many graphs: e.g. if Ψ is compatible with γ and if $\gamma' \geq \gamma$, then Ψ is compatible with γ' as well, which follows from (3.3).

If the function $\psi : \mathcal{A}_\gamma \rightarrow \mathbb{C}$ in (3.4) is smooth we say that the corresponding cylindrical function $\Psi : \mathcal{A} \rightarrow \mathbb{C}$ is smooth as well. Let

$$\text{Cyl}^\infty := \text{span} \{ \Psi \mid \Psi \text{ is a smooth cylindrical function on } \mathcal{A} \}.$$

In fact, Cyl^∞ is not only a linear space, but an algebra also. To see this, note first that every element Ψ' of the space is a finite linear combination of smooth cylindrical functions $\{\Psi_i\}$ ($i = 1, \dots, k$) compatible, respectively, with graphs $\{\gamma_i\}$. There exists a graph γ such that $\gamma \geq \gamma_i$ for every i under consideration which means that every function Ψ_i is compatible with the graph. Hence Ψ' is a smooth cylindrical function compatible with γ also. Now, given elements Ψ and Ψ' of Cyl^∞ we can find a graph γ' such that the two functions are compatible with it. Then $\Psi\Psi'$ is a smooth cylindrical function compatible with γ' , thus an element of Cyl^∞ .

3.2 Flux operators

Let us now define some differential operators on Cyl^∞ . Let S be a $(d-1)$ -dimensional regular analytic manifold with fixed (external) orientation and P_S be the restriction of the bundle P to S . Denote by λ a smooth map $\mathbb{R} \times P_S \rightarrow P_S$ such that for each arbitrary but fixed τ the map $\lambda_\tau := \lambda(\tau, \cdot)$ is a vertical automorphism of P_S such that:

$$\lambda_{\tau+\tau'} = \lambda_\tau \circ \lambda_{\tau'} \text{ and } \lambda_0 = \text{id},$$

i.e. λ_τ is a one-parameter group of vertical automorphisms of P_S .

To define the differential operators on Cyl^∞ we distinguish graphs *adapted* to S —we say that a graph γ is adapted to S iff every edge of γ either: (i) is contained in S (modulo its endpoints), or (ii) does not intersect S at all, or (iii) intersects S at exactly one endpoint. Next, among edges belonging to the class (iii) we distinguish ones placed ‘up’ (‘down’) the *oriented* submanifold S . Note that if a graph γ' is not adapted to S , then, thanks to the analyticity of S and the edges of γ' , one can adapt the graph to the submanifold by the appropriate subdividing of its edges i.e. for every γ' there exists $\gamma \geq \gamma'$ such that γ is adapted to S [6]. Thus every cylindrical function is compatible with a graph adapted to a given S .

Suppose now that a graph γ is adapted to S . By means of a given family λ_τ of vertical automorphisms of P_S we define a family of smooth

maps³ $\lambda_\tau : \mathcal{A}_\gamma \rightarrow \mathcal{A}_\gamma$ by the formula

$$\lambda_\tau(A_\gamma) := (\lambda_\tau^{I_t} A(e_I) \lambda_\tau^{I_s}), \quad (3.5)$$

where $\lambda_\tau^{I_s}$ and $\lambda_\tau^{I_t}$ are automorphisms of the fibres $P_{(e_I)_s}$ and $P_{(e_I)_t}$ respectively defined as follows:

1. if the edge e_I belongs to the class (i) or (ii) described above, then $\lambda_\tau^{I_t} = \text{id}$ and $\lambda_\tau^{I_s} = \text{id}$;
2. if the edge e_I belongs to the class (iii) and
 - (a) $S \cap e_I = (e_I)_s$, i.e. e_I is 'outgoing' from S then $\lambda_\tau^{I_s}$ acts on the fibre $P_{(e_I)_s}$ as
 - i. λ_τ if e_I is placed 'up' the submanifold S ,
 - ii. $\lambda_{-\tau}$ if e_I is placed 'down' the submanifold S ,
and $\lambda_\tau^{I_t} = \text{id}$.
 - (b) if $S \cap e_I = (e_I)_t$ i.e. e_I is 'ingoing' to S then $\lambda_\tau^{I_t}$ acts on the fibre $P_{(e_I)_t}$ as
 - i. $\lambda_{-\tau}$ if e_I is placed 'up' the submanifold S ,
 - ii. λ_τ if e_I is placed 'down' the submanifold S ,
and $\lambda_\tau^{I_s} = \text{id}$.

Let $\Psi = p_\gamma^* \psi$ be a smooth cylindrical function compatible with the graph γ adapted to S .

Definition 3.4 *Flux operator $\hat{X}_{S,\lambda}$ associated with the submanifold S and the family λ_τ of vertical automorphisms acts on Ψ in the following way:*

$$\hat{X}_{S,\lambda} \Psi := -\frac{i}{2} \frac{d}{d\tau} \Big|_{\tau=0} (p_\gamma^* \lambda_\tau^* \psi).$$

Although the function Ψ is compatible with many distinct graphs adapted to S the r.h.s. of the above equation does not depend on the choice of an adapted graph. Clearly, $\hat{X}_{S,\lambda}$ is a linear operator defined on the whole Cyl^∞ and preserves the space. It is easy to see that in the case of a trivial bundle

³The maps λ_τ ($\tau \in \mathbb{R}$) define smooth curves on $\mathcal{A}_\gamma$, which cannot be lifted to curves on $\mathcal{A}$. But as it was noted by Professor S.L. Woronowicz the maps do define curves on the space $\overline{\mathcal{A}}$ of generalized connections (for the definition of $\overline{\mathcal{A}}$ see Footnote 5).

Definition (3.4) is equivalent to the standard definition of flux operators introduced in [6].

For S, S_1, S_2 such that $S = S_1 \cup S_2$ and $S_1 \cap S_2 = \emptyset$ and for any λ_τ defined on P_S we have:

$$\hat{X}_{S,\lambda} = \hat{X}_{S_1,\lambda_1} + \hat{X}_{S_2,\lambda_2}, \quad (3.6)$$

where $(\lambda_i)_\tau := \lambda_\tau|_{P_{S_i}}$ ($i = 1, 2$).

3.3 Definition of the Sahlmann algebra

Let $\mathfrak{A}$ be a complex algebra generated by all operators on Cyl^∞ of the following form:

$$\begin{aligned} \Psi &\mapsto \hat{\Phi}\Psi := \Phi\Psi, \\ \Psi &\mapsto \hat{X}_{S,\lambda}\Psi, \end{aligned} \quad (3.7)$$

where $\Psi, \Phi \in \text{Cyl}^\infty$. Define $*$ operation on $\mathfrak{A}$:

$$\hat{\Phi}^* := \hat{\Phi}, \quad \hat{X}_{S,\lambda}^* := \hat{X}_{S,\lambda}.$$

To show that the $*$ operation is well defined on $\mathfrak{A}$ note first that the above formulas define a $*$ operation on the free algebra $\mathfrak{A}_F$ generated by all the smooth cylindrical functions and the flux operators. Obviously, every element a of $\mathfrak{A}_F$ defines a linear operator on Cyl^∞ via (3.7), which will be denoted by a also. Then

$$\mathfrak{A} = \mathfrak{A}_F / \mathfrak{A}_0,$$

where $\mathfrak{A}_0$ is a (left and right) ideal in $\mathfrak{A}_F$ defined as follows:

$$\mathfrak{A}_0 := \{a \in \mathfrak{A}_F \mid a\Psi = 0 \text{ for all } \Psi \in \text{Cyl}^\infty\}.$$

Now, the $*$ operation is well defined on $\mathfrak{A}$ if

$$a \in \mathfrak{A}_0 \implies a^* \in \mathfrak{A}_0. \quad (3.8)$$

To prove the implication fix an element a of $\mathfrak{A}_F$ and a smooth cylindrical function Ψ . The element a is generated by a finite number of smooth cylindrical functions and flux operators. It is possible to find a graph γ such that (i) γ is adapted to all the surfaces corresponding to the flux operators, (ii) all the cylindrical functions are compatible with γ and (iii) the function Ψ is compatible with γ . Now it is easy to see that $a\Psi$ is also compatible with γ .

Consider now the space $\mathcal{A}_\gamma$. As it was mentioned earlier, any trivialization of the bundle $P_{V(\gamma)}$ defines a bijection between $\mathcal{A}_\gamma$ and G^N , where N is the number of the edges of γ . This bijection can be used to push-forward the Haar measure on G^N onto $\mathcal{A}_\gamma$ [10]. The resulting measure $d\mu_\gamma$ is independent of the choice of the trivialization of $P_{V(\gamma)}$ and can be used to define a scalar product $\langle \cdot | \cdot \rangle_\gamma$ on the linear space of smooth cylindrical functions compatible with the graph γ . Then we have [6]

$$\langle \Psi | a \Psi' \rangle_\gamma = \langle a^* \Psi | \Psi' \rangle_\gamma$$

for every pair Ψ, Ψ' of smooth cylindrical functions compatible with γ . Thus if $a \in \mathfrak{A}_0$ then

$$\langle a^* \Psi | \Psi' \rangle_\gamma = 0.$$

Taking into account smoothness of $a^* \Psi$ we conclude that $a^* \Psi = 0$, which proves the implication (3.8).

Now we are able to give a definition of the Sahlman algebra:

Definition 3.5 *Sahlmann holonomy-flux *-algebra (Sahlmann algebra for short) is the *-algebra $(\mathfrak{A}, *)$ [1, 2].*

We emphasize that the Sahlmann algebra is a unital one with a unit given by a constant cylindrical function of the value equal to 1.

4 Induced action of automorphisms on $\mathfrak{A}$

There is a natural (right) action of the group of automorphisms of P on the space $\mathcal{A}$:

$$\mathcal{A} \ni A \mapsto \theta^* A \in \mathcal{A}, \quad (4.1)$$

where $\theta \in \Theta$. Given $\theta \in \Theta^\omega$ and $A \in \mathcal{A}$ the parallel transport $(\theta^* A)(e) : P_{e_s} \rightarrow P_{e_t}$ can be expressed as

$$(\theta^* A)(e) = \text{ad}_{\theta^{-1}}(A(\check{\theta}(e))). \quad (4.2)$$

This suggests to define a map⁴:

$$\mathcal{A}_\gamma \ni A_\gamma \mapsto \text{Ad}_\theta A_\gamma := (\text{ad}_\theta (A(e_I))) \in \mathcal{A}_{\check{\theta}(\gamma)}.$$

⁴The map Ad_θ is an analytic diffeomorphism, what can be easily seen after expressing the map in any trivializations of bundles $P_{V(\gamma)}$ and $P_{V(\check{\theta}(\gamma))}$.

where edges $\{e_I\}$ constitute the graph γ . Applying (4.2) to the edges we obtain

$$p_\gamma \circ \theta^* = \text{Ad}_{\theta^{-1}} \circ p_{\tilde{\theta}(\gamma)},$$

where the maps on both sides of the latter equation are from $\mathcal{A}$ onto $\mathcal{A}_\gamma$.

The right action (4.1) induces a linear left action of θ (that is, a representation of the group Θ) on the space of functions on $\mathcal{A}$:

$$(\tilde{\theta}\Psi)(A) := \Psi(\theta^*A),$$

where Ψ is a function on $\mathcal{A}$. Let Ψ_n be a sequence of functions on $\mathcal{A}$ converging pointwisely to a function Ψ . Then

$$\lim_{n \rightarrow \infty} \tilde{\theta}\Psi_n = \tilde{\theta}(\lim_{n \rightarrow \infty} \Psi_n) = \tilde{\theta}\Psi, \quad (4.3)$$

which means that $\tilde{\theta}$ is continuous with respect to the topology of pointwise convergence.

We have the following lemma:

Lemma 4.1 *If $\theta \in \Theta^\omega$ and $\Psi \in \text{Cyl}^\infty$ then $\tilde{\theta}\Psi \in \text{Cyl}^\infty$.*

Proof. Let $\Psi \in \text{Cyl}^\infty$ be compatible with the graph γ . Then

$$\begin{aligned} (\tilde{\theta}\Psi)(A) &= \Psi(\theta^*A) = \psi(p_\gamma(\theta^*A)) = \psi(\text{Ad}_{\theta^{-1}} p_{\tilde{\theta}(\gamma)}(A)) = \\ &= [p_{\tilde{\theta}(\gamma)}^*(\text{Ad}_{\theta^{-1}}^* \psi)](A), \end{aligned}$$

that is:

$$\tilde{\theta}\Psi = p_{\tilde{\theta}(\gamma)}^*(\text{Ad}_{\theta^{-1}}^* \psi), \quad (4.4)$$

where $\text{Ad}_{\theta^{-1}}^* \psi$ is a smooth complex function on $\mathcal{A}_{\tilde{\theta}(\gamma)}$. Thus $\tilde{\theta}\Psi$ is a smooth cylindrical function compatible with the (analytic) graph $\tilde{\theta}(\gamma)$. ■

Now we can define the action of the automorphism θ on the Sahlmann algebra:

$$\mathfrak{A} \ni \hat{a} \mapsto \tilde{\theta}\hat{a}\tilde{\theta}^{-1}. \quad (4.5)$$

Lemma 4.2 *The map (4.5) is a $*$ -preserving isomorphism of the algebra $\mathfrak{A}$ onto itself.*

The above lemma is an implication of the following one:

Lemma 4.3 (i) If $\Phi \in \text{Cyl}^\infty$, then:

$$\tilde{\theta} \hat{\Phi} \tilde{\theta}^{-1} = \widehat{\tilde{\theta} \Phi}.$$

(ii) For flux operators the following formula is true:

$$\tilde{\theta} \hat{X}_{S,\lambda} \tilde{\theta}^{-1} = \hat{X}_{\tilde{\theta}(S), \text{ad}_\theta \lambda},$$

where $(\text{ad}_\theta \lambda)_\tau := \text{ad}_\theta \lambda_\tau$.

Proof of the statement (ii) (the proof of (i) is trivial). We have:

$$\begin{aligned} \frac{2}{i} \tilde{\theta} (\hat{X}_{S,\lambda} \Psi) &= \tilde{\theta} \left[\frac{d}{d\tau} \Big|_{\tau=0} (p_\gamma^* \lambda_\tau^* \psi) \right] = \\ &= \frac{d}{d\tau} \Big|_{\tau=0} [\tilde{\theta} (p_\gamma^* \lambda_\tau^* \psi)] = \frac{d}{d\tau} \Big|_{\tau=0} [p_{\tilde{\theta}(\gamma)}^* (\text{Ad}_{\tilde{\theta}^{-1}}^* \lambda_\tau^* \psi)] = \\ &= \frac{d}{d\tau} \Big|_{\tau=0} [p_{\tilde{\theta}(\gamma)}^* (\text{Ad}_{\tilde{\theta}^{-1}}^* \lambda_\tau^* \text{Ad}_\theta^*) (\text{Ad}_{\tilde{\theta}^{-1}}^* \psi)]. \quad (4.6) \end{aligned}$$

In the second step of above calculation we used the linearity of $\tilde{\theta}$ and (4.3) to change the order of the action of $\tilde{\theta}$ and the differentiation with respect to τ , in the third one we applied (4.4) to the function $p_\gamma^*(\lambda_\tau^* \psi)$ as cylindrical one compatible with the graph γ .

Definition (3.5) and Equation (2.1) imply that the family of maps $\text{Ad}_\theta \circ \lambda_\tau \circ \text{Ad}_{\tilde{\theta}^{-1}}$ acting on (and preserving) the space $\mathcal{A}_{\tilde{\theta}(\gamma)}$ is defined by the family $(\text{ad}_\theta \lambda)_\tau$ of vertical automorphisms of $P_{\tilde{\theta}(S)}$, i.e.

$$\text{Ad}_\theta \circ \lambda_\tau \circ \text{Ad}_{\tilde{\theta}^{-1}} = (\mathbf{ad}_\theta \lambda)_\tau.$$

Setting this to the r.h.s of Equation (4.6) we get:

$$\tilde{\theta} (\hat{X}_{S,\lambda} \Psi) = \hat{X}_{\tilde{\theta}(S), \text{ad}_\theta \lambda} (\tilde{\theta} \Psi).$$

■

5 Representations of Sahlmann algebra

The Sahlmann algebra $\mathfrak{A}$ is not equipped with any norm. This does not allow us to expect that one can represent the algebra by means of *bounded* operators on a Hilbert space. The fact leads to the following definition of a $*$ -representation of $\mathfrak{A}$ [2]:

Definition 5.1 Let $L(\mathcal{H})$ be a space of linear operators on a Hilbert space $\mathcal{H}$. We say that a map $\pi : \mathfrak{A} \rightarrow L(\mathcal{H})$ is a $*$ -representation of $\mathfrak{A}$ on the Hilbert space $\mathcal{H}$ if:

1. there exists a dense subspace $\mathcal{D}$ of $\mathcal{H}$ such that

$$\mathcal{D} \subset \bigcap_{\hat{a} \in \mathfrak{A}} [D(\pi(\hat{a})) \cap D(\pi^*(\hat{a}))],$$

where $D(\pi(\hat{a}))$ denotes the domain of the operator $\pi(\hat{a})$;

2. for every $\hat{a}, \hat{b} \in \mathfrak{A}$ and $\lambda \in \mathbb{C}$ the following conditions are satisfied on $\mathcal{D}$:

$$\begin{aligned} \pi(\hat{a} + \hat{b}) &= \pi(\hat{a}) + \pi(\hat{b}), & \pi(\lambda \hat{a}) &= \lambda \pi(\hat{a}), \\ \pi(\hat{a} \hat{b}) &= \pi(\hat{a}) \pi(\hat{b}), & \pi(\hat{a}^*) &= \pi^*(\hat{a}). \end{aligned}$$

3. If, given $\hat{a} \in \mathfrak{A}$, there exists a subspace E dense in $\mathcal{H}$ such that $E \subset D(\pi(\hat{a}))$ and $\pi(\hat{a})|_E$ is closable, then $\pi(\hat{a})$ is equal to the closure of $\pi(\hat{a})|_E$.

We say that π is non-degenerate iff the fact that for every $\hat{a} \in \mathfrak{A}$ $\pi(\hat{a})v = 0$ implies $v = 0$.

These conditions mean in particular that $\pi(\hat{a})\mathcal{D} \subset \mathcal{D}$ and that every element $\hat{a} = \hat{a}^*$ is represented by the operator $\pi(\hat{a})$ symmetric on $\mathcal{D}$.

In the sequel we will consider only non-degenerate $*$ -representations of $\mathfrak{A}$.

5.1 Basic facts

Although Sahlman algebra is not equipped with any norm we can define a norm on its subalgebra Cyl^∞ :

$$\|\hat{\Psi}\|_{\text{sup}} := \sup_{A \in \mathcal{A}} |\Psi(A)|.$$

If π is a non-degenerate $*$ -representation of $\mathfrak{A}$ on $\mathcal{H}$, then $\pi|_{\text{Cyl}^\infty}$ maps elements of Cyl^∞ into *bounded* operators on $\mathcal{H}$ and is uniquely extendable to a (non-degenerate) representation of a C^* -algebra Cyl defined as the completion of Cyl^∞ in the norm $\|\cdot\|_{\text{sup}}$ [2]. The connection between the representations $\pi|_{\text{Cyl}^\infty}$ and the representations of the C^* -algebra Cyl implies the following characterization of π and $\mathcal{H}$ [1]:

Characterization 5.1 (Sahlmann)

1. The representation $\pi|_{\text{Cyl}^\infty}$ is a direct sum of cyclic representations $\{\pi_\nu\}$:

$$\mathcal{H} = \bigoplus_{\nu \in \mathcal{N}} \mathcal{H}_\nu, \quad \pi|_{\text{Cyl}^\infty} = \bigoplus_{\nu \in \mathcal{N}} \pi_\nu, \quad (5.1)$$

where $\{\mathcal{H}_\nu\}$ are carrier spaces of representations $\{\pi_\nu\}$, respectively, ν ranges some label set $\mathcal{N}$ and the sum is orthogonal;

2. For each ν there exists a Hilbert space isomorphism

$$\varphi_\nu : L^2(\overline{\mathcal{A}}, \mu_\nu) \rightarrow \mathcal{H}_\nu, \quad (5.2)$$

where $\overline{\mathcal{A}}$ is a Gel'fand-Neimark spectrum⁵ of Cyl , and μ_ν is a regular, Borel measure on $\overline{\mathcal{A}}$.

Every Hilbert space $\mathcal{H}_\nu$ is defined by the choice of a vector $v_\nu \in \mathcal{H}$:

$$\mathcal{H}_\nu := \overline{\{\pi(\hat{\Phi})v_\nu \mid \Phi \in \text{Cyl}^\infty\}}, \quad (5.3)$$

the measure μ_ν is given by the following formula:

$$\int_{\overline{\mathcal{A}}} \Phi d\mu_\nu := \langle v_\nu | \pi(\hat{\Phi})v_\nu \rangle, \quad (5.4)$$

and the isomorphism φ_ν is the closure of the map:

$$L^2(\overline{\mathcal{A}}, \mu_\nu) \supset \text{Cyl}^\infty \ni \Phi \mapsto \pi(\hat{\Phi})v_\nu \in \mathcal{H}_\nu \quad (5.5)$$

(here $\langle \cdot | \cdot \rangle$ is the scalar product on $\mathcal{H}$). Clearly, the set of isomorphisms $\{\varphi_\nu\}$ defines an isomorphism φ , which maps $\bigoplus_{\nu \in \mathcal{N}} L^2(\overline{\mathcal{A}}, \mu_\nu)$ onto $\mathcal{H}$. In the sequel we will assume that φ is fixed and will identify both the Hilbert spaces. Similarly, we will not distinguish between representation π on $\mathcal{H}$ and $\varphi^{-1}\pi\varphi$ on $\bigoplus_{\nu \in \mathcal{N}} L^2(\overline{\mathcal{A}}, \mu_\nu)$ and both will be denoted by π .

⁵There exists the following geometric characterization of the spectrum of Cyl . Let $B_{x,y}$ ($x, y \in \Sigma$) be a set of all morphisms from P_x onto P_y . The set $B := \bigcup B_{x,y}$, where (x, y) runs over $\Sigma \times \Sigma$, possesses a natural groupoid structure. A *generalized connection* $\bar{A}$ on P is a homomorphism from the groupoid of analytic oriented edges in Σ into B such that it maps every oriented edge e to a morphism $\bar{A}(e) \in B_{(e)_s, (e)_t}$. The fact that $\bar{A}$ is a homomorphism means that it satisfies conditions (3.1). The spectrum of Cyl can be identified with the set $\overline{\mathcal{A}}$ of all generalized connections on P . Clearly, $\mathcal{A} \subset \overline{\mathcal{A}}$. For more details see [9].

Given $\Psi \in \mathcal{H}$, we will denote by Ψ_ν the orthogonal projection of Ψ onto $\mathcal{H}_\nu = L^2(\overline{\mathcal{A}}, \mu_\nu)$ and write $\Psi = (\Psi_\nu)$. Following [1] we define

$$\mathcal{C}^\infty := \{(\Psi_\nu) \in \mathcal{H} \mid \Psi_\nu = 0 \text{ for all but finitely many } \nu\text{'s} \\ \text{and } \Psi_\nu \in \text{Cyl}^\infty \text{ for every } \nu\}. \quad (5.6)$$

Since Cyl^∞ is dense in $L^2(\overline{\mathcal{A}}, \mu_\nu)$, $\mathcal{C}^\infty$ is dense in $\mathcal{H}$. It turns out that the assumption $\mathcal{D} = \mathcal{C}^\infty$, where $\mathcal{D}$ is introduced by Definition 5.1, gives the following useful characterization [1]:

Characterization 5.2 (Sahlmann) *Suppose π is a non-degenerate $*$ -representation of $\mathfrak{A}$ in a Hilbert space $\mathcal{H}$, and π satisfies the assumption $\mathcal{D} = \mathcal{C}^\infty$. For every $\hat{X}_{S,\lambda}$, π defines a family of elements of $\mathcal{H}$ labelled by elements of the set $\mathcal{N}$ (the same as in Equation (5.1))*

$$\mathcal{N} \ni \iota \mapsto F_{S,\lambda}^\iota \in \mathcal{H}, \quad (5.7)$$

such that the following conditions are satisfied (given $F_{S,\lambda}^\iota$, we will subsequently denote by $F_{S,\lambda}^\iota{}_\nu$ the $\mathcal{H}_\nu$ component of $F_{S,\lambda}^\iota$):

1. for every $\Psi = (\Psi_\nu) \in \mathcal{C}^\infty$:

$$\pi(\hat{X}_{S,\lambda})\Psi = \hat{\mathbf{X}}_{S,\lambda}\Psi + \hat{F}_{S,\lambda}\Psi$$

where $\hat{\mathbf{X}}_{S,\lambda}\Psi := (\hat{X}_{S,\lambda}\Psi_\nu)$ and:

$$\hat{F}_{S,\lambda}\Psi = \hat{F}_{S,\lambda}(\Psi_\nu) := \left(\sum_\iota \Psi_\iota F_{S,\lambda}^\iota{}_\nu\right),$$

where $\sum_\iota \Psi_\iota F_{S,\lambda}^\iota{}_\nu$ belongs to $\mathcal{H}_\nu$.

2. for every $\Phi, \Phi' \in \text{Cyl}^\infty \subset \mathcal{H}_\nu$

$$\langle \hat{X}_{S,\lambda}\Phi | \Phi' \rangle_\nu - \langle \Phi | \hat{X}_{S,\lambda}\Phi' \rangle_\nu = \langle \Phi | (F_{S,\lambda}^\nu{}_\nu - \overline{F}_{S,\lambda}^\nu{}_\nu) \Phi' \rangle_\nu, \quad (5.8)$$

where $\langle \cdot | \cdot \rangle_\nu$ is the scalar product on $\mathcal{H}_\nu$ (**no** summation with respect to the index $\nu \in \mathcal{N}$ in $F_{S,\lambda}^\nu{}_\nu$);

3. for every $S = S_1 \cup S_2$ such that S_i ($i = 1, 2$) are disjoint:

$$F_{S,\lambda}^\iota{}_\nu = F_{S_1,\lambda_1}^\iota{}_\nu + F_{S_2,\lambda_2}^\iota{}_\nu, \quad (5.9)$$

where $(\lambda_i)_\tau := \lambda_\tau|_{P_{S_i}}$.

5.2 Automorphism covariant representations of Sahlmann algebra

Following [2] we formulate:

Definition 5.2 *Suppose π is a non-degenerate $*$ -representation of the Sahlmann algebra $\mathfrak{A}$ on a Hilbert space $\mathcal{H}$. We will say that π is an automorphism covariant representation if and only if there exists a family $\{v_\nu\}$ ($\nu \in \mathcal{N}$) of vectors in $\mathcal{H}$ such that (i) $\mathcal{H}$ admits the orthogonal decomposition $\mathcal{H} = \bigoplus_{\nu \in \mathcal{N}} \mathcal{H}_\nu$, where each $\mathcal{H}_\nu$ is defined by v_ν according to (5.3) and (ii) every finite linear combination v of v_ν 's defines an automorphism invariant state on $\mathfrak{A}$:*

$$\langle v | \pi(\tilde{\theta} \hat{a} \tilde{\theta}^{-1}) v \rangle = \langle v | \pi(\hat{a}) v \rangle \quad (5.10)$$

for every $a \in \mathfrak{A}$, and every automorphism $\theta \in \Theta^\omega$.

The consequences of the above definition are the following. Vectors $\{v_\nu\}$ define an isomorphism φ , which identifies spaces $\bigoplus_{\nu \in \mathcal{N}} L^2(\overline{\mathcal{A}}, \mu_\nu)$ and $\mathcal{H}$. One can easily check that in terms of this identification the linear subspace $\mathcal{D} := \mathcal{C}^\infty$ of $\mathcal{H}$ satisfies the requirements of the Definition 5.1. This means that Characterization 5.2 is applicable to every automorphism covariant $*$ -representation of the Sahlmann algebra.

The requirement (5.10) implies that the map

$$\pi(\hat{\Phi})v_\nu \mapsto \pi(\tilde{\theta} \hat{\Phi} \tilde{\theta}^{-1})v_\nu,$$

(where $\Phi \in \text{Cyl}^\infty$) defined on $\mathcal{H}_\nu$ is closable and its closure u_θ is a unitary map on $\mathcal{H}_\nu$. This allow us to conclude that the measure μ_ν on $\overline{\mathcal{A}}$ defined by (5.4) is *automorphism invariant*, i.e.:

$$\int_{\overline{\mathcal{A}}} \Psi d\mu_\nu = \int_{\overline{\mathcal{A}}} (u_\theta \Psi) d\mu_\nu. \quad (5.11)$$

Because of the unitarity of u_θ the map

$$\mathcal{H} \ni \Psi \mapsto U_\theta \Psi := (u_\theta \Psi_\nu) \in \mathcal{H}$$

defines a unitary representation of Θ^ω on $\mathcal{H}$:

$$\theta \mapsto U_\theta.$$

Finally, for every $\hat{a} \in \mathfrak{A}$ the equation

$$\pi(\tilde{\theta} \hat{a} \tilde{\theta}^{-1}) = U_\theta \pi(\hat{a}) U_\theta^{-1}$$

is satisfied on $\mathcal{D}$. Indeed, given $\theta \in \Theta^\omega$, every element v of $\mathcal{D}$ can be written as

$$v = \sum_{\nu} \pi(\tilde{\theta} \hat{\Psi}_{\nu} \tilde{\theta}^{-1}) v_{\nu}.$$

Hence for any $v, v' \in \mathcal{D}$

$$\begin{aligned} \langle v | \pi(\tilde{\theta} \hat{a} \tilde{\theta}^{-1}) v' \rangle &= \sum_{\nu \nu'} \langle \pi(\tilde{\theta} \hat{\Psi}_{\nu} \tilde{\theta}^{-1}) v_{\nu} | \pi(\tilde{\theta} \hat{a} \tilde{\theta}^{-1}) \pi(\tilde{\theta} \hat{\Psi}'_{\nu'} \tilde{\theta}^{-1}) v_{\nu'} \rangle = \\ &= \sum_{\nu \nu'} \langle v_{\nu} | \pi(\tilde{\theta} \hat{\Psi}_{\nu}^* \hat{a} \hat{\Psi}'_{\nu'} \tilde{\theta}^{-1}) v_{\nu'} \rangle = \sum_{\nu \nu'} \langle \pi(\hat{\Psi}_{\nu}) v_{\nu} | \pi(\hat{a}) \pi(\hat{\Psi}'_{\nu'}) v_{\nu'} \rangle = \\ &= \langle v | U_{\theta} \pi(\hat{a}) U_{\theta}^{-1} v' \rangle \end{aligned}$$

On $\mathcal{D}$ we also have (see Lemma 4.3):

$$U_{\theta} \hat{\mathbf{X}}_{S, \lambda} U_{\theta}^{-1} = \hat{\mathbf{X}}_{\tilde{\theta}(S), \text{ad}_{\theta} \lambda}, \quad U_{\theta} \hat{F}_{S, \lambda} U_{\theta}^{-1} = \hat{F}_{\tilde{\theta}(S), \text{ad}_{\theta} \lambda}. \quad (5.12)$$

6 Main theorem

Given a $*$ -representation π of the Sahlmann algebra $\mathfrak{A}$ on a Hilbert space $\mathcal{H}$, every decomposition (5.1) of $\mathcal{H}$ is determined by a choice of the vectors $\{v_{\nu}\}$ ($\nu \in \mathcal{N}$) defining spaces $\mathcal{H}_{\nu}$ according to (5.3). On the other hand, the representation π and the vectors $\{v_{\nu}\}$ define the family $\{\mu_{\nu}\}$ of measures on the space of generalized connection $\overline{\mathcal{A}}$ by means of the formula (5.4).

Theorem 6.1 *Suppose π is a non-degenerate, automorphism covariant $*$ -representation of the Sahlmann algebra $\mathfrak{A}$. Then for every member of the family $\{v_{\nu}\}$ ($\nu \in \mathcal{N}$) of vectors satisfying the requirements of Definition 5.2 the corresponding measure*

$$\mu_{\nu} = \mu_{\text{AL}},$$

where μ_{AL} is a (defined below) natural measure on $\overline{\mathcal{A}}$. In the consequence, all the $L^2(\overline{\mathcal{A}}, \mu_{\text{AL}})$ functions $F_{S, \lambda}^{\nu}$ (no summation) used in Characterization 5.2 are real valued.

Remarks

1. The theorem is a generalization of the theorem 5.1 in [2]. The difference between the two theorems is that in [2] we assumed $\Sigma = \mathbb{R}^d$ and $P = \Sigma \times G$, while the present theorem is valid for a Sahlmann algebra built on an arbitrary principal bundle $P(\Sigma, G)$ of a compact connected structure group G . Moreover, in [2] we used the notion of a *diffeomorphism* covariant representation of the Sahlmann algebra, which is not natural while working with an arbitrary (non-trivial) bundle $P(\Sigma, G)$.

2. Upon the requirement of non-degeneracy and the definition of the automorphism covariance, every $*$ -representation π of the Sahlmann $*$ -algebra satisfies the conclusions of the theorem.

To define μ_{AL} let us recall that any trivialization of a bundle $P_{V(\gamma)}$ defines a bijection between $\mathcal{A}_\gamma$ and G^N , where N is the number of the edges of the graph γ . The bijection allows us to push forward the (normalized) Haar measure from G^N onto $\mathcal{A}_\gamma$. Denote the resulting measure by μ_γ . It is easy to show that μ_γ does not depend on the choice of the trivialization of the bundle $P_{V(\gamma)}$. Let $\psi : \mathcal{A}_\gamma \rightarrow \mathbb{C}$ be a continuous function. The measure μ_{AL} is the only measure on $\overline{\mathcal{A}}$ satisfying the requirement [10]

$$\int_{\overline{\mathcal{A}}} (\bar{p}_\gamma^* \psi) d\mu_{\text{AL}} = \int_{\mathcal{A}_\gamma} \psi d\mu_\gamma,$$

where $\bar{p}_\gamma : \overline{\mathcal{A}} \rightarrow \mathcal{A}_\gamma$ is a projection defined analogously⁶ to the projection p_γ (see Subsection 3.1).

7 Proof of the main theorem

We have already concluded that Characterizations 5.1 and 5.2 are applicable to every $*$ -representation π satisfying the assumptions of the main theorem. According to the latter characterization the operator $\pi(\hat{X}_{S,\lambda})$ has the following form on $\mathcal{C}^\infty$:

$$\pi(\hat{X}_{S,\lambda}) = \hat{\mathbf{X}}_{S,\lambda} + \hat{F}_{S,\lambda}. \quad (7.1)$$

It turns out that the automorphism covariance of the representation imposes a restriction on the operator $\hat{F}_{S,\lambda}$ in (7.1)—we will show that there exists a submanifold C and a one-parameter group Λ_τ of vertical automorphisms of P_C such that for every $\nu \in \mathcal{N}$ the imaginary part of $F_{C,\Lambda}^\nu$ is equal to zero. This fact and Equation (5.8) imply that the operator $\hat{X}_{C,\Lambda}$ is *symmetric* on $\text{Cyl}^\infty \subset L^2(\overline{\mathcal{A}}, \mu_\nu)$. Then we will show that the set of symmetric flux operators is rich enough to use Lemma 6.3 of [2] to conclude that $\mu_\nu = \mu_{\text{AL}}$ for every $\nu \in \mathcal{N}$.

⁶The projection $\bar{p}_\gamma$ is defined by an equivalence relation on $\overline{\mathcal{A}}$ analogous to the relation (3.2) on $\mathcal{A}$. Denote by $\bar{A}_\gamma$ the equivalence class of a generalized connection $\bar{A}$ i.e., $\bar{A}_\gamma =: \bar{p}_\gamma(\bar{A})$. Consequently we denote $\overline{\mathcal{A}}_\gamma := \bar{p}_\gamma(\overline{\mathcal{A}})$. In fact, every equivalence class $\bar{A}_\gamma$ contains a smooth connection $A \in \mathcal{A}$ [10], hence $\overline{\mathcal{A}}_\gamma$ is naturally isomorphic to $\mathcal{A}_\gamma$. This justifies the expression $\bar{p}_\gamma : \overline{\mathcal{A}} \rightarrow \mathcal{A}_\gamma$ used above.

7.1 The imaginary part of $F_{C,\Lambda}^\nu$

7.1.1 The functions $F_{S,\lambda}^\ell$

Consider the functions $F_{S,\lambda}^\ell$ and their imaginary parts:

$$I_{S,\lambda}^\ell := \frac{1}{2}(F_{S,\lambda}^\ell - \overline{F}_{S,\lambda}^\ell) \in L^2(\overline{\mathcal{A}}, \mu_\nu).$$

The assumed automorphism covariance of the representation π allows us to make use of the results derived in Subsection 5.2. Equation (5.12) implies

$$U_\theta \hat{F}_{S,\lambda} \Psi = \hat{F}_{\check{\theta}(S), \text{ad}_\theta \lambda} U_\theta \Psi,$$

for every automorphism $\theta \in \Theta^\omega$ and for every $\Psi \in \mathcal{C}^\infty$. This means that the elements $F_{S,\lambda}^\ell$ of $\mathcal{H}$ are assigned to the submanifolds S and to the family λ_τ in a covariant way,

$$u_\theta F_{S,\lambda}^\ell = F_{\check{\theta}(S), \text{ad}_\theta \lambda}^\ell, \quad \text{hence} \quad u_\theta I_{S,\lambda}^\ell = I_{\check{\theta}(S), \text{ad}_\theta \lambda}^\ell. \quad (7.2)$$

The automorphism covariance of the representation π also implies that, for every ν the scalar product $\langle \cdot | \cdot \rangle_\nu$ on $L^2(\overline{\mathcal{A}}, \mu_\nu)$ is automorphism invariant, thus in particular

$$\|I_{S,\lambda}^\ell\|_\nu = \|u_\theta I_{S,\lambda}^\ell\|_\nu = \|I_{\check{\theta}(S), \text{ad}_\theta \lambda}^\ell\|_\nu. \quad (7.3)$$

Equation (5.9) allows us to conclude that for $S = S_1 \cup S_2$, where S_1 and S_2 are disjoint

$$I_{S,\lambda}^\ell = I_{S_1, \lambda_1}^\ell + I_{S_2, \lambda_2}^\ell \quad (7.4)$$

(here $(\lambda_i)_\tau := \lambda_\tau|_{P_{S_i}}$ and $i = 1, 2$).

7.1.2 The functions $I_{C,\Lambda}^\nu$

Let us fix the index ν and denote $I_{S,\lambda} = I_{S,\lambda}^\nu$ in order to simplify notation. Recall that $I_{S,\lambda} \in L^2(\overline{\mathcal{A}}, \mu_\nu)$.

Consider a quintuplet

$$(C, c, \chi, \omega_s, \lambda_\tau)$$

where C, c are $(d-1)$ -dimensional submanifolds⁷ of Σ , χ is an element of Θ^ω , ω_s ($s \in \mathbb{R}$) is a family of automorphisms belonging to Θ^ω and λ_τ is

⁷Precisely speaking we require C and c to be $(d-1)$ -dimensional *manifolds with boundary* embedded in Σ . In the sequel of the proof we will associate with C and c some flux operators. Note that although we have defined flux operators only for submanifolds the definition can be extended in a natural way to involve more general objects—consider submanifolds $S' \subset S$ and $\lambda'_\tau = \lambda_\tau|_{P_{S'}}$. Then $\hat{X}_{S,\lambda} - \hat{X}_{S',\lambda'}$ is a flux operator associated with $S \setminus S'$ being an embedded manifold with boundary. This justifies the existence of the flux operators associated with C and c .

a one-parameter group of vertical automorphism of P_c , (where P_c is the restriction of P to c). Assume that the members of the quintuplet satisfy the following conditions:

1. $C \cup c = \check{\chi}(c)$ and $C \cap c = \emptyset$;
2. every $\check{\omega}_s$ preserves c , i.e. for every $s \in \mathbb{R}$

$$\check{\omega}_s(c) = c;$$

3. for every point $y \in C$ there exists $s_y \in \mathbb{R}$ such that $s > s_y$ implies $\check{\omega}_s(y) \notin C$.
4. automorphisms χ and ω_s preserve the family λ_τ i.e. for every $\tau, s \in \mathbb{R}$ the following vertical automorphisms on P_c are equal to each other:

$$(\text{ad}_\chi \lambda_\tau)|_{P_c} = \lambda_\tau = \text{ad}_{\omega_s} \lambda_\tau \quad (7.5)$$

Define now the following family of vertical automorphisms of P_C :

$$\Lambda_\tau := (\text{ad}_\chi \lambda_\tau)|_{P_C}. \quad (7.6)$$

We are going to find a relation between $I_{c,\lambda}$ and $I_{C,\Lambda}$. Equation (7.3) gives:

$$\|I_{\check{\chi}(c), \text{ad}_\chi \lambda}\|_\nu^2 = \|I_{c,\lambda}\|_\nu^2.$$

$\check{\chi}(c) = C \cup c$ and $c \cap C = \emptyset$, thus by virtue of Equation (7.4):

$$I_{\check{\chi}(c), \text{ad}_\chi \lambda} = I_{c,\lambda} + I_{C,\Lambda}$$

(note that in the above equation we used the first equality of Equation (7.5)). Combining the two latter equations we obtain:

$$\|I_{C,\Lambda}\|_\nu^2 = -2\langle I_{c,\lambda} | I_{C,\Lambda} \rangle_\nu. \quad (7.7)$$

Our goal now is to show that the scalar product $\langle I_{c,\lambda} | I_{C,\Lambda} \rangle_\nu$ is equal to 0. We will do it in two steps: first, we will express the function $I_{c,\lambda}$ as a limit of a suitably chosen sequence of cylindrical functions belonging to $\text{Cyl}^\infty \subset L^2(\overline{\mathcal{A}}, \mu_\nu)$. Then, Characterization 5.2 will allow us to conclude that $\langle I_{c,\lambda} | I_{C,\Lambda} \rangle_\nu = 0$.

7.1.3 The function $I_{c,\lambda}$ as a limit of a sequence of cylindrical functions

The function $I_{c,\lambda}$ can be expressed as a limit,

$$I_{c,\lambda} = \lim_{n \rightarrow \infty} \Phi_n; \quad \Phi_n \in \text{Cyl}^\infty.$$

Note that, in fact, we have quite a large freedom in the choice of the sequence converging to $I_{c,\lambda}$ —to see this, consider a sequence of automorphisms (θ_n) such that (i) every $\check{\theta}_n$ is analytic and preserves the submanifold c and (ii) every θ_n preserves the family λ_τ (i.e. $\text{ad}_{\theta_n} \lambda_\tau = \lambda_\tau$). Then, applying (7.2) and the unitarity of u_θ we obtain:

$$\|I_{c,\lambda} - \Phi_n\|_\nu = \|u_{\theta_n}(I_{c,\lambda} - \Phi_n)\|_\nu = \|I_{c,\lambda} - u_{\theta_n}(\Phi_n)\|_\nu,$$

which means that $\lim_{n \rightarrow \infty} u_{\theta_n}(\Phi_n) = I_{c,\lambda}$ as well. We will use this freedom to construct some special sequence, which converges to $I_{c,\lambda}$.

Let us fix n and consider a graph γ_n compatible with the cylindrical function Φ_n . In general, some edges of the graph can be transversal⁸ to the submanifold C . Then the action of $\hat{X}_{C,\Lambda}$ on Φ_n is (in general) nontrivial,

$$\hat{X}_{C,\Lambda} \Phi_n \neq 0.$$

Lemma 7.1 *For each of the graphs γ_n , $n = 1, 2, \dots$ defined above, there exists an automorphism $\theta_n \in \Theta^\omega$, such that:*

1. $\check{\theta}_n$ preserves the submanifold c and the action ad_{θ_n} preserves the family λ_τ ;
2. the graph $\check{\theta}_n(\gamma_n)$ has no edges transversal to the submanifold C .

Proof. Fix n and denote by $\{y_1, \dots, y_M\}$ the set of the intersection points between C and edges of γ_n transversal to C . The properties of the quintuplet $(C, c, \chi, \omega_s, \lambda_\tau)$ imply the existence of a set $\{s_1, \dots, s_M\}$ of real numbers such that:

$$\tilde{\omega}_{s_J}(y_J) \notin C.$$

⁸An edge e is *transversal* to a submanifold S iff $S \cap e$ is one-element set. Without loss of generality C can be expressed as $S \setminus S'$ for some submanifolds S and S' . Now, an edge e is by definition *transversal* to C if e is transversal to S and $S \cap e = C \cap e$. An appropriate subdivision of each edge transversal to C gives edges of the class (iii) (Subsection 3.2) with respect to submanifold S . Thus only transversal edges can give non-zero terms contributing to the action of $\hat{X}_{C,\Lambda}$.

Let $s(n) := \max\{s_1, \dots, s_M\}$. Then

$$\theta_n := \omega_{s(n)}$$

preserves c and the family λ_τ and thus satisfies the requirements of the lemma. ■

Use now the automorphisms (θ_n) given by Lemma 7.1 to construct the following sequence convergent to $I_{c,\lambda}$,

$$\tilde{\Phi}_n := u_{\theta_n}(\Phi_n).$$

Now, each function in the sequence $(\tilde{\Phi}_n)$ is compatible with a graph having no edge transversal to the submanifold C , hence

$$\hat{X}_{C,\Lambda} \tilde{\Phi}_n = 0. \quad (7.8)$$

7.1.4 The vanishing of $\langle I_{c,\lambda} | I_{C,\Lambda} \rangle_\nu$

Equation (5.8) in the case $\Phi' = 1$ gives

$$\langle \hat{X}_{C,\Lambda} \Phi | 1 \rangle_\nu = 2 \langle \Phi | I_{C,\Lambda} \rangle_\nu,$$

for every smooth cylindrical function $\Phi \in L^2(\overline{\mathcal{A}}, \mu_\nu)$. Owing to Equation (7.8)

$$0 = \lim_{n \rightarrow \infty} \langle \hat{X}_{C,\Lambda} \tilde{\Phi}_n | 1 \rangle_\nu = \lim_{n \rightarrow \infty} 2 \langle \tilde{\Phi}_n | I_{C,\Lambda} \rangle_\nu = 2 \langle I_{c,\lambda} | I_{C,\Lambda} \rangle_\nu.$$

This result and Equation (7.7) imply

$$\|I_{C,\Lambda}\|_\nu^2 = 0, \quad \text{hence} \quad I_{C,\Lambda} = I_{C,\Lambda}^\nu = 0, \quad (7.9)$$

where the last equality refers to elements of $L^2(\overline{\mathcal{A}}, \mu_\nu)$ while the measure μ_ν is not assumed to be faithful.

7.2 Construction of the quintuplet

To justify the result (7.9) we have to show that there exists a quintuplet $(C, c, \chi, \omega_s, \lambda_\tau)$ satisfying all the requirements described in Subsection 7.1.2. This will be done by an explicit construction.

To construct the quintuplet we have to take into account that the automorphisms χ and ω_s have to be elements of Θ^ω i.e., that they have to define *globally* analytic diffeomorphisms $\tilde{\chi}$ and $\tilde{\omega}_s$ on Σ . On the other hand, the

diffeomorphisms have to transform (and, in particular, preserve) submanifolds C and c in the required way. Because it is rather difficult to construct an analytic diffeomorphism which e.g. preserves a given submanifold we will first construct the diffeomorphisms and only then we will find appropriate submanifolds C and c .

The idea of the construction of the quintuplet is the following: (i) first we will find an analytic vector field Y on Σ such that it generates one-parameter group of analytic diffeomorphisms on Σ ; (ii) the diffeomorphisms $\tilde{\chi}$ and $\tilde{\omega}_s$ will be defined as maps which shift points along the integral curves of Y , i.e. preserve the curves; (iii) the submanifolds C and c will be constructed by the shift of a $(d-2)$ -dimensional submanifold along the integral curves of Y ; (iv) we will obtain automorphisms χ and ω_s by a horizontal lift (with respect to a connection on P) of $\tilde{\chi}$ and $\tilde{\omega}_s$; (v) finally we will use a section of the bundle P_c to define λ_τ .

7.2.1 A construction of analytic diffeomorphisms

It is well known that every (sufficiently regular) vector field on a manifold generates a *local* one-parameter group of *local* diffeomorphisms on this manifold (see e.g. [14]), which in the case of non-compact manifolds is not always extendable to a group of diffeomorphisms. Therefore to obtain (a one-parameter group of) diffeomorphisms on Σ we have to choose the generating vector field in a suitable way. To choose it we will apply a theorem by Grauert [15], which states that every analytic manifold can be embedded into $\mathbb{R}^k$ of sufficiently high dimension by means of an *analytic proper* map. Because the properness of the map will be crucial for the below considerations let us recall the definition of the notion.

Let T_1, T_2 be topological spaces. A continuous map $\xi : T_1 \rightarrow T_2$ is proper if and only if for every compact $U \subset T_2$ the inverse image $\xi^{-1}(U)$ is compact in T_1 . One can show in particular that $\xi(T_1)$ is closed in T_2 and if ξ is injective, then ξ is homeomorphism between T_1 and $\xi(T_1)$ with topology induced by the topology of T_2 (see e.g. [16]).

Let us assume then that Σ is embedded into some $\mathbb{R}^k$ by an analytic proper map. Denote by q the Riemannian metric on Σ induced by the Euclidean metric on $\mathbb{R}^k$. Then the following lemma holds:

Lemma 7.2 *Suppose Y is an analytic vector field on Σ such that the function $q(Y, Y)$ is bounded on Σ . Then Y generates a one-parameter group $\tilde{\alpha}_t$ of analytic diffeomorphisms on Σ .*

Before we prove the lemma let us note that the lemma is not true for an arbitrary Riemannian metric on Σ , as it is shown by the following example:

Example 7.1 Let $\Sigma = \mathbb{R}$ and $g(\partial_x, \partial_x) = \exp(-x^2)$, then for $Y = x^2 \partial_x$ function $g(Y, Y)$ is bounded on $\mathbb{R}$, but Y does not generate a group of diffeomorphisms on $\mathbb{R}$. ■

Proof of Lemma 7.2. Every analytic vector field generates a local one-parameter group of local *analytic* diffeomorphisms (see e.g. [17]). Thus to prove the lemma it is enough to show that every *non-extendable* integral curve of Y is a map from the *whole* $\mathbb{R}$ into Σ . Let us consider then a non-extendable integral curve:

$$\mathbb{R} \supset]a, b[\ni t \mapsto y(t) \in \Sigma. \quad (7.10)$$

where a, b can be equal to $\pm\infty$ respectively.

The theory of ODE's (see e.g. [17]) guarantees that if $Y(y(t_0)) = 0$ for some $t_0 \in]a, b[$, then $y(t) \equiv y(t_0)$ and consequently $]a, b[= \mathbb{R}$. If a given integrable curve forms a loop, then obviously: $]a, b[= \mathbb{R}$.

Now let us suppose that $Y(y(t)) \neq 0$ on $]a, b[$ and the integral curve does not form a loop. Parametrize the path $y([a, b])$ by its length τ defined by the metric g . Thus after reparametrization of the curve $y(t)$ we get a curve:

$$\mathbb{R} \supset]a', b'[\ni \tau \mapsto y'(\tau) \in \Sigma. \quad (7.11)$$

We have:

$$Y(y'(\tau)) = \frac{d\tau}{dt} \partial_\tau, \quad \text{hence} \quad q(Y, \partial_\tau) = \frac{d\tau}{dt} > 0$$

(here ∂_τ is a unital vector field on the submanifold $y([a, b])$ generated by the curve $y'(\tau)$.) Integrating the latter equation we obtain:

$$t(\tau) = \int_{\tau_0}^{\tau} \frac{d\tau'}{q(Y, \partial_\tau)}, \quad (7.12)$$

where τ_0 is chosen in a way satisfying $t(a') = a$ and $t(b') = b$.

Assume that $b' = \infty$ ($a' = -\infty$). Because $q(Y, \partial_\tau) > 0$ and is bounded (by virtue of the Schwarz inequality and the boundedness of $q(Y, Y)$) the l.h.s. of Equation (7.12) is not bounded from above (below), hence $b = \infty$ ($a = -\infty$).

Suppose now that $b' < \infty$ and $b = t(b') < \infty$. Then:

$$\lim_{t \rightarrow b} y(t) = \lim_{\tau \rightarrow b'} y'(\tau) = \mathbf{y} \in \Sigma. \quad (7.13)$$

Actually, the existence of the limit (7.13) is a consequence of the fact that the metric q is induced by the Euclidean metric⁹ on $\mathbb{R}^k$, hence we can consider the curve $y'(\tau)$ as a curve in $\mathbb{R}^k$ and find the limit in this space. Because Σ is a closed subset of $\mathbb{R}^k$ (which follows from the properness of the map embedding Σ in the $\mathbb{R}^k$) the limit belongs to Σ .

The theory of ODE's applied to the *analytic* vector field Y ensures that there exists a neighbourhood $U \subset \Sigma$ of $\mathbf{y}$ and $\delta > 0$ such that for every $(t_0, y_0) \in]-\delta, \delta[\times U$ there exists *exactly one* curve $\tilde{y} :]-\delta, \delta[\rightarrow \Sigma$ being an integral curve of Y and satisfying the initial condition $\tilde{y}(t_0) = y_0$. The assumption $b < \infty$ implies that

1. if $Y(\mathbf{y}) \neq 0$ then the curve $y(t)$ under consideration is extendable, which contradicts our assumption about the curve,
2. If $Y(\mathbf{y}) = 0$ then there exist two *distinct* solutions of ODE $dy/ds = Y$ with the initial value $y(0) = \mathbf{y}$: one of them is $y(s) \equiv \mathbf{y}$ and the second one is:

$$y(s) = \begin{cases} y(s) = y'(\tau(s)) & \text{for } s < 0 \\ y(s) = \mathbf{y} & \text{for } s \geq 0 \end{cases},$$

where $\tau(s)$ is given by $\int_{\tau(s)}^{b'} d\tau/q(Y, \partial_\tau) = -s$. Clearly, this is forbidden by the theory of ODE's as written above.

In this way we conclude that $b = \infty$. Similarly we have $a = -\infty$ even if $a' > -\infty$. Thus $]a, b[= \mathbb{R}$. ■

Example 7.2 Let us fix a point $y_0 \in \Sigma$ and a nonzero vector $Y_{y_0} \in T_{y_0}\Sigma$ and suppose for simplicity that Σ is embedded into some $\mathbb{R}^k$ in such a way that y_0 is placed at the origin of the Cartesian coordinate frame of $\mathbb{R}^k$. Clearly, Y_{y_0} defines a constant vector field on $\mathbb{R}^k$. Let $\tilde{Y}$ be a vector field on $\mathbb{R}^k$ given by the following formula:

$$\tilde{Y}(x) := \exp(-\|x\|_{\mathbb{R}}^2) Y_{y_0},$$

where $x \in \mathbb{R}^k$ and $\|\cdot\|_{\mathbb{R}}$ is the standard norm on $\mathbb{R}^k$. Let Y be a vector field on Σ defined as the orthogonal (with respect to the standard metric of $\mathbb{R}^k$) projection of $\tilde{Y}$ onto the tangent bundle $T\Sigma$. Then Y is an analytic vector field on Σ satisfying the assumption of Lemma 7.2 (and condition $Y(y_0) = Y_{y_0}$). Hence Y generates a one-parameter group $\tilde{\alpha}_t$ of analytic diffeomorphisms on Σ . ■

⁹If q was an arbitrary metric on Σ , then $\lim_{\tau \rightarrow b'} y'(\tau)$ could not exist — consider Σ and g as in Example 7.1. Then $\Sigma = \mathbb{R}$ can be parametrized by length parameter $]a', b'[\ni \tau \mapsto y'(\tau) \in \Sigma$, where a', b' are finite. Clearly $\lim_{\tau \rightarrow b'} y'(\tau) = \infty$.

In this way we finished the step (i) of the construction.

7.2.2 Construction of the quadruplet $(C, c, \check{\chi}, \check{\omega}_s)$

Now let us construct the quadruplet $(C, c, \check{\chi}, \check{\omega}_s)$ satisfying the first three conditions described in Subsection 7.1.2 (steps (ii) and (iii) of the construction).

Fix an arbitrary point $y_0 \in \Sigma$ and consider the following ingredients, from which the quadruplet $(C, c, \check{\chi}, \check{\omega}_s)$ will be built:

1. Let Y be an analytic vector field on Σ satisfying assumption of Lemma 7.2 such that $Y(y_0) \neq 0$. Denote by $\check{\alpha}_t$ the one-parameter group of diffeomorphisms generated by Y .
2. Let σ be a $(d-1)$ -dimensional analytic (possibly non-closed) submanifold of Σ such that $y_0 \in \sigma$ and $Y(y_0)$ is *not* tangent to σ .
3. Let $\xi : \Sigma \rightarrow \mathbb{R}$ be a bounded analytic function such that for some neighbourhood $\tilde{U}$ of y_0 (i) $\xi = 0$ on $\tilde{U} \cap \sigma$ and (ii) $\xi > 0$ on $\tilde{U} \setminus \sigma$ (such a function can be easily constructed by means of the Grauert theorem — see Example 7.3).

Then there exists a neighbourhood $u \subset \sigma$ of point y_0 , an (analytic) coordinate frame $(y^1, \dots, y^{d-1})$ on u and $t_0 > 0$ such that:

1. (t, y^i) is an analytic coordinate frame on:

$$U := \{\check{\alpha}_t(u) \mid t \in]-3t_0, 3t_0[\} \subset \tilde{U} \subset \Sigma$$

2. for every $y \in U$:

$$Y(y) \neq 0. \tag{7.14}$$

Now let us define a function ξ' as the pull-back of ξ :

$$\xi' := \check{\alpha}_{t_0}^* \xi$$

and the two vector fields satisfying assumption of Lemma 7.2 as

$$Z := \xi' Y \quad \text{and} \quad W := \xi \xi' Y. \tag{7.15}$$

Denote by $\check{\beta}_s$ and $\check{\omega}_s$ the one-parameter group of diffeomorphisms generated by the vector fields Z and W respectively.




Figure 1: Construction of submanifolds C and c

Consider now sets (see Figure 1):

$$\begin{aligned}\check{\alpha}(u) &:= \{\check{\alpha}_t(u) \mid t \in]-t_0, t_0] \} \\ \check{\alpha}^+(u) &:= \{\check{\alpha}_t(u) \mid t \in]0, t_0] \} \\ \check{\alpha}^-(u) &:= \{\check{\alpha}_t(u) \mid t \in]-t_0, 0] \}\end{aligned}$$

(we emphasize that $\check{\alpha}^-(u)$ contains u ; clearly $\check{\alpha}(u) = \check{\alpha}^+(u) \cup \check{\alpha}^-(u)$). The properties of the function ξ guarantee that ξ' is *non-zero* on $\check{\alpha}(u)$ and that it is equal to zero on $\check{\alpha}_{-t_0}(u)$. Taking into account Equation (7.14), one can easily conclude that if $s \neq 0$, the diffeomorphism $\check{\beta}_s$ is not the identity on the whole $\check{\alpha}(u)$ and it is the identity on $\check{\alpha}_{-t_0}(u)$ for every $s \in \mathbb{R}$. Similarly, for $s \neq 0$ diffeomorphism $\check{\omega}_s$ is not the identity on $\check{\alpha}(u) \setminus u$ and is the identity on $\check{\alpha}_{-t_0}(u)$ and u .

Except for the points where vector fields Z and W vanish, the integral curves of Y, Z and W define the same paths. Thus every $\check{\omega}_s$ preserves $\check{\alpha}^-(u)$. Moreover, for every point $y \in \check{\alpha}^+(u)$ there exists $s_y \in \mathbb{R}$ such that $s > s_y$ implies¹⁰ $\check{\omega}_s(y) \notin \check{\alpha}^+(u)$.

¹⁰Note that $\check{\alpha}^+(u)$ does *not* contain any points, at which $\check{\omega}_{s \neq 0}$ is the identity. Moreover, u is the only subset of the boundary of $\check{\alpha}^+(u)$, where $\check{\omega}_s = \text{id}$. Thus if s_y is large enough

Now we are ready to define the quadruplet $(C, c, \check{\chi}, \check{\omega}_s)$. In fact, $\check{\omega}_s$ is already defined as a family of diffeomorphisms generated by W . It is possible to choose (i) $(d-2)$ -dimensional regular analytic submanifold¹¹ c_u contained in u and containing the point y_0 and (ii) a positive number s_0 such that the following analytic $(d-1)$ -dimensional submanifold

$$C := \{\check{\beta}_s(c_u) \mid s \in]0, s_0]\}$$

is contained in $\check{\alpha}^+(u)$. Define:

$$c := \{\check{\alpha}_t(c_u) \mid t \in]-t_0, 0]\}$$

(clearly, c is contained in $\check{\alpha}^-(u)$) and:

$$\check{\chi} := \check{\beta}_{s_0}. \quad (7.16)$$

Hence we have $C \cup c = \check{\chi}(c)$ and $C \cap c = \emptyset$. In this way we completed the construction of the quadruplet $(C, c, \check{\chi}, \check{\omega}_s)$ satisfying the first three requirements described in Subsection 7.1.2.

Example 7.3 (continuation of Example 7.2) To construct a quadruplet $(C, c, \check{\chi}, \check{\omega}_s)$ using the vector field Y described in Example 7.2 we only need to define the submanifold σ and the function ξ . Let $(\cdot|\cdot)$ be the standard scalar product on $\mathbb{R}^k$. Define the function

$$\mathbb{R}^k \ni x \mapsto \tilde{\xi}(x) := 1 - \exp[-(x|Y(y_0))^2] \in \mathbb{R}.$$

Clearly $\tilde{\xi}$ is bounded nonnegative function on $\mathbb{R}^k$ and it vanishes on the $(k-1)$ -dimensional plane $\mathbb{P}$ orthogonal to the vector $Y(y_0)$. Let $B(r, y_0)$ be a ball in $\mathbb{R}^k$ of center in y_0 and radius $r > 0$. Thus we can define:

$$\begin{aligned} \xi &:= \tilde{\xi}|_{\Sigma} \\ \sigma &:= \mathbb{P} \cap B(r_0, y_0) \cap \Sigma \end{aligned}$$

for some (small enough) $r_0 > 0$. ■

$\check{\omega}_{s_y}$ shifts a given point $y \in \check{\alpha}^+(u)$ out this set. Note as well that because $\check{\omega}_s$ is the identity on u the integral curves of W passing through $\check{\alpha}^+(u)$ do not form loops—hence $\check{\omega}_s(y) \notin \check{\alpha}^+(u)$, only if $s > s_y$.

¹¹If $\dim \Sigma = 2$ take $c_u = y_0$.

7.2.3 Construction of automorphisms χ , ω_s and λ_τ

Now, by suitable lifts of diffeomorphisms $\check{\chi}$ and $\check{\omega}_s$ we will get the desired automorphisms χ and ω_s (step (iv)) and we will then define the family λ_τ of vertical automorphisms on P_c (step (v)), which will complet the construction of the quintuplet $(C, c, \chi, \omega_s, \lambda_\tau)$.

Let T denote a neighbourhood of the point y_0 (around which submanifolds C and c were just constructed) such that the map

$$\text{pr}^{-1}(T) \ni p \mapsto \varsigma(p) = (\text{pr}(p), g_p) \in T \times G$$

is a local trivialization of the bundle P . It is clear that we can construct the manifolds C, c in such a way that $c \subset T$. Moreover, without loss of generality we can assume that there exist neighbourhoods T_0, T_1 of y_0 such that (i) $c \subset T_0 \subset T_1 \subset T$ and (ii) there exists a smooth function η on Σ equal to 0 on T_0 and 1 on $\Sigma \setminus T_1$.

Let A be any connection one-form on P . Define the connection A_0 in the following way:

$$\begin{cases} A_0 := A & \text{on } \text{pr}^{-1}(\Sigma \setminus T) \\ \varsigma_0^* A_0 := \eta \varsigma_0^* A & \text{on } T \end{cases}$$

where ς_0 is a local section of P over T defined as $\varsigma(\varsigma_0(y)) = (y, \mathbb{I})$ ($y \in T$, $\mathbb{I}$ is the neutral element of G). Note that on T_0 the pull-back $\varsigma_0^* A_0 = 0$, which means that every set $\varsigma^{-1}(y, g = \text{const})$, $y \in T_0$ defines a (local) section of P which is *horizontal* with respect to the connection A_0 .

Let Z^* and W^* be defined by A_0 horizontal lifts of the vector fields Z and W (see Equation (7.15)), respectively. Denote by β_s and ω_s the one-parameter groups of diffeomorphisms on P generated by Z^* and W^* , respectively. Clearly, the diffeomorphisms β_s and ω_s are automorphisms of P and elements of Θ^ω . Define now:

$$\chi := \beta_{s_0},$$

where s_0 is the same real number as in (7.16).

Let f be a constant function from c to the Lie algebra G' . The desired family λ_τ can be defined using the trivialization ς in the following way:

$$P_c \ni p \mapsto \lambda_\tau(p) := (\text{pr}(p), \exp(\tau f)g_p) \in P_c. \quad (7.17)$$

In the trivialization ς the automorphisms χ and ω_s act in the following way:

$$\chi(p) = (\check{\chi}(\text{pr}(p)), g_p), \quad \omega_s(p) = (\check{\omega}_s(\text{pr}(p)), g_p),$$

provided $p \in \text{pr}^{-1}(T_0)$ —this is because, by virtue of the construction, the vector fields Z^* and W^* are tangent to the submanifolds defined by $(\text{pr}(p), g_p = \text{const})$. Thus it is easy to see that λ_τ satisfies the condition (7.5).

This completes the construction of the quintuplet $(C, c, \chi, \omega_s, \lambda_\tau)$.

7.3 Final conclusion

Lemma 6.2 of [2] can be reformulated in the following way (see also [18]):

Lemma 7.3 *Let G be a compact, connected Lie group and μ be a probability (regular Borel) measure on $\overline{\mathcal{A}}$. Suppose that for every collection of edges $\{e_1, \dots, e_N\}$ constituting a graph there exists $N \times \dim G$ operators X_{S_I, λ_I^i} ($I = 1, \dots, N$, $i = 1, \dots, \dim G$) such that (i) the operators X_{S_I, λ_I^i} are symmetric on $\text{Cyl}^\infty \subset L^2(\overline{\mathcal{A}}, \mu)$ (ii) for every I the intersection $S_I \cap (e_1 \cup \dots \cup e_N) = S_I \cap e_I$ is a one-element set and e_I is transversal to S_I (iii) for every I the families $\{\lambda_I^i\}$ generate a reper field on $P_{S_I \cap e_I}$. Then:*

$$\mu = \mu_{\text{AL}}.$$

Before we show that in our case all assumptions of the lemma are satisfied let us make some remarks. The lemma means that for every $\nu \in \mathcal{N}$ the measure $\mu_\nu = \mu_{\text{AL}}$, thus it completes the proof of the first conclusion of the main theorem. On the other hand, *every* operator $\hat{X}_{S, \lambda}$ is self-adjoint on $L^2(\overline{\mathcal{A}}, \mu_{\text{AL}})$ [6]. Thus the first conclusion and Equation (5.8) imply the second conclusion of the theorem i.e. that for every ν the function $F_{S, \lambda}^\nu$ is real valued.

Let us turn back to the assumptions of the lemma. Equation (5.8) and (7.9) imply that for C and Λ_τ considered in the last subsection and for every smooth cylindrical functions Φ, Φ' :

$$\langle \hat{X}_{C, \Lambda} \Phi | \Phi' \rangle_\nu - \langle \Phi | \hat{X}_{C, \Lambda} \Phi' \rangle_\nu = 0,$$

which means that $\hat{X}_{C, \Lambda}$ is *symmetric* on $\text{Cyl}^\infty \subset L^2(\overline{\mathcal{A}}, \mu_\nu)$. Moreover, for every $\theta \in \Theta^\omega$ the operator $\hat{X}_{\tilde{\theta}(C), \text{ad}_\theta \Lambda}$ is symmetric on Cyl^∞ as well. Therefore now we only have to show that there exist sufficiently many operators of the form $\hat{X}_{\tilde{\theta}(C), \text{ad}_\theta \Lambda}$, i.e. that the assumptions (ii) and (iii) are satisfied by the operators.

Note that for every edge $e \subset \Sigma$ there exists a submanifold C (being a member of a quintuplet) and an automorphism $\theta \in \Theta^\omega$ such that a submanifold $\tilde{\theta}(C)$ intersects e at exactly one point not being any vertex of e (this is

required in fact by the assumption (ii)) and e is transversal to the submanifold. To see this, choose a point $y_0 \in \text{Int } e$ and a vector $Y_{y_0} \in T_{y_0} \Sigma$ such that Y_{y_0} is *not* tangent to e . Applying the construction described in Examples 7.2 and 7.3 we obtain a vector field Y with a family of diffeomorphisms $\tilde{\alpha}_t$ and a quintuplet containing a submanifold C —this submanifold can be defined (by an appropriate choice of $(d-2)$ -dimensional submanifold c_u) in such a way that $\tilde{\alpha}_{-t_0}(C)$ for some (small enough) $t_0 > 0$ meets e transversally at exactly one interior point of the edge. Then one can find $\theta \in \Theta^\omega$ such that $\tilde{\theta} = \tilde{\alpha}_{-t_0}$. Thus e is transversal to $\tilde{\theta}(C)$ and the assumption (ii) is satisfied.

Consider now the submanifold c constructed together with the submanifold C . It is easy to choose $n = \dim G$ families $\{\lambda_\tau^i\}$ of vertical automorphisms acting on P_c such that they generate a reper field on every fibre of the bundle¹²—indeed, we have constructed the family λ_τ by means of the (constant) function¹³ $f : c \rightarrow G'$ (Equation (7.17)), therefore n constant functions $\{f^i\}$ define desired families $\{\lambda_\tau^i\}$ provided the values of these functions form a basis of G' . Thus the families

$$\text{ad}_\theta \Lambda_\tau^i = \text{ad}_\theta[(\text{ad}_\chi \lambda_\tau^i)|_{P_C}]$$

generate (in particular) a reper field on the fibre over the intersection point between $\tilde{\theta}(C)$ and the edge e . This means that the assumption (iii) is satisfied.

In this way we have finished the proof of the main theorem.

8 Summary

In this paper we considered automorphism covariant $*$ -representations of the Sahlmann holonomy-flux algebra $(\mathfrak{A}, *)$ for a theory of connections on an arbitrary bundle $P(\Sigma, G)$, where Σ is an arbitrary real-analytic manifold and G is a compact connected Lie group. We showed that the carrier space of the $*$ -representations used in Sahlmann's Characterization 5.2 is the orthogonal product (Theorem 6.1):

$$\mathcal{H} = \bigoplus_{\nu} L^2(\overline{\mathcal{A}}, \mu_\nu), \quad (8.1)$$

where every measure μ_ν is the natural measure:

$$\mu_\nu = \mu_{\text{AL}}. \quad (8.2)$$

¹²Here the notion of a reper field concerns the fibre thought as a separate manifold.

¹³ G' indicates the Lie algebra of the group G .

We emphasize that, although the requirement of the automorphism covariance of the representation singles out the measure μ_{AL} , there may exist inequivalent automorphism covariant representations of the Sahlmann algebra on the Hilbert space (8.1) which differ from each other by the family of functions $F_{S,\lambda}{}^\ell{}_\nu$ (note that the functions are not fully determined by Theorem 6.1). Thus automorphism covariance with respect to the group of automorphisms whose projections are *analytic* diffeomorphisms, (i.e. the group Θ^ω) does not seem to fix a unique automorphism covariant representation.

It is, however, possible to define an action of a larger group of automorphisms on the holonomy-flux $*$ -algebra and modify slightly the definition of the automorphism covariance of representations of the algebra. Then all the considerations presented here and Theorem 6.1 remain valid. We can also go a step further and prove that the automorphism covariance with respect to the extended group of automorphisms singles out precisely one representation of the Sahlmann algebra which is the cyclic representation described by Characterization 5.2 with all the correction terms equal zero. This result will be presented in [12].

Acknowledgements: Our interest in this subject was stimulated by discussions with Hanno Sahlmann who also explained to us many technical points and discussed with us his unpublished considerations. Stanisław L. Woronowicz encouraged us to reformulate our results in the language of non-trivial principal bundles. We are grateful to Marcin Bobieński for a discussion which helped us to formulate Lemma 6.2. Finally, we thank Abhay Ashtekar and Thomas Thiemann for numerous discussions about the quantum geometry and, respectively, QSD. This work was supported in part by the KBN grants 2 PO3B 068 23 and 2 PO3B 127 24 and by the NSF grant PHY 0090091.

References

- [1] Sahlmann H 2002 Some Comments on Representation Theory of the Algebra Underlying Loop Quantum Gravity *Preprint* gr-qc/0207111
- [2] Okołów A and Lewandowski J 2003 Diffeomorphism covariant representations of the holonomy-flux $*$ -algebra *Class. Quant. Grav.* **20**, 3543–3567 *Preprint* gr-qc/0302059

- [3] Thomas Thiemann 2004 *Modern Canonical Quantum General Relativity* (Cambridge University Press) at press *Preprint* gr-qc/0110034
C. Rovelli 2004 *Quantum Gravity* (Cambridge University Press, Cambridge)
Ashtekar A and Lewandowski J 2004 Background Independent Quantum Gravity: A Status Report *Preprint* gr-qc/0404018
- [4] Ashtekar A, Lewandowski J, Marolf D, Mourão J and Thiemann T 1995 Quantization of diffeomorphism invariant theories of connections with local degrees of freedom *J. Math. Phys.* **36** 6456–6493, *Preprint* gr-qc/9504018
- [5] Ashtekar A and Isham C J 1992 Representations of the holonomy algebras for gravity and non-Abelian gauge theories *Class. Quant. Grav.* **9** 1433–1467
- [6] Ashtekar A and Lewandowski J 1997 Quantum theory of geometry. I: Area operators *Class. Quant. Grav.* **14** A55–A82 *Preprint* gr-qc/9602046
- [7] Ashtekar A, Corichi A and Zapata J A 1998 Quantum Theory of Geometry III: Non-commutativity of Riemannian Structures *Class. Quant. Grav.* **15** 2955–2972 *Preprint* gr-qc/9806041
- [8] Sahlmann H 2002 When Do Measures on the Space of Connections Support the Triad Operators of Loop Quantum Gravity? *Preprint* gr-qc/0207112
- [9] Ashtekar A and Lewandowski J 1995 Differential geometry on the space of connections using projective techniques, *Jour. Geo. & Phys.* **17**, 191–230
- [10] Ashtekar A and Lewandowski J 1994 Representation theory of analytic holonomy C^* -algebras *Knots and quantum gravity* (Baez J (ed), Oxford: Oxford University Press) *Preprint* gr-qc/9311010
Marolf D and Mourão J 1995 On the support of the Ashtekar-Lewandowski measure *Commun. Math. Phys.* **170**, 583–606.
Ashtekar A and Lewandowski J 1995 Projective techniques and functional integration for gauge theories *J. Math. Phys.* **36** 2170–2191 *Preprint* gr-qc/9411046
- [11] Sahlmann H and Thiemann T 2003 On the Superselection Theory of the Weyl Algebra for Diffeomorphism Invariant Quantum Gauge Theories *Preprint* gr-qc/0302090

- [12] Lewandowski J, Okołów A, Sahlmann H and Thiemann T Uniqueness of diffeomorphism invariant representation of holonomy-flux algebra (in preparation)
- [13] Baez J C 1996 Spin Network States in Gauge Theory *Adv. Math.* **117** 253–272 *Preprint* gr-qc/9411007
- [14] Kobayashi S and Nomizu K 1963 *Foundations of Differential Geometry* (New York; London: Interscience Publisher)
- [15] Grauert H 1958 On Levi's Problem and the Imbedding of Real-Analytic Manifolds *Ann. Math.* **68** 460–472
- [16] Bourbaki N 1989 *General topology* (Berlin; New York: Springer-Verlag)
- [17] Maurin K 1976 *Analysis* (Dordrecht; Boston: D. Reidel Pub. Co.)
- [18] Okołów A 2004 Representations of Quantum Geometry, Ph.D. Thesis, University of Warsaw